

Spontaneous CP Violation at Finite Temperature in the MSSM¹

Koichi Funakubo^{a,2}, Akira Kakuto^{b,3},
Shoichiro Otsuki^{b,4} and Fumihiko Toyoda^{b,5}

^{a)}*Department of Physics, Saga University, Saga 8408502 Japan*

^{b)}*Kyushu School of Engineering, Kinki University, Iizuka 8208255 Japan*

By studying the effective potential of the MSSM at finite temperature, we find that CP can be spontaneously broken in the intermediate region between the symmetric and broken phases separated by the bubble wall created at the phase transition. If CP is violated in this manner, there could be a bubble wall connecting CP conserving vacua and violating CP halfway, which would result in sufficient baryon asymmetry of the universe.

1 Introduction

Within the framework of electroweak theory, the baryon asymmetry of the universe (BAU) could be explained if the electroweak phase transition (EWPT) in the early universe is of first order[1]. The minimal standard model with sufficiently heavy Higgs boson ($m_H > 67\text{GeV}$) is found to have second-order EWPT. Further the effect of the only CP violation in the KM matrix is too weak to generate the present BAU. Hence, one is led to extend the standard model in such a way that the extra source of CP violation exists and the bosonic content is enlarged to have the EWPT stronger. Among such extensions, the minimal supersymmetric standard model (MSSM) with the small soft-supersymmetry-breaking parameter in the stop mass-squared matrix[2] will cause the first-order EWPT and offer many sources of CP violation. The CP violations effective for generation of the chiral charge flux at the tree level are the relative phases among the μ -parameter, the gaugino mass parameters and the A -parameters included in the mass matrices of the charginos, neutralinos and the scalar partners of the quarks and leptons[3]. Besides these phases, the relative phase of the two Higgs doublets not only enters

¹Talk by K. Funakubo at the Workshop on Fermion Mass and CP Violation, Hiroshima, Japan, March 1998.

²e-mail: funakubo@cc.saga-u.ac.jp

³e-mail: kakuto@fuk.kindai.ac.jp

⁴e-mail: otkslscp@mbox.nc.kyushu-u.ac.jp

⁵e-mail: ftoyoda@fuk.kindai.ac.jp

these mass matrices but also directly couple to the quarks and leptons to yield the flux carried by these particles into the symmetric phase region[4, 5]. In general, the phases of the Higgs scalars vary spatially around the bubble wall so that the gauge-invariant relative phase cannot be rotated away from the Yukawa interactions.

In a previous paper[6] we attempted to determine the profile of the bubble wall by solving the equations of motion for the effective potential at the transition temperature ($> T_C \simeq 100\text{GeV}$) in the two-Higgs-doublet model. For some set of parameters, we presented a solution such that CP -violating phase spontaneously generated becomes as large as $O(1)$ around the wall while it completely vanishes in the broken and symmetric phase limits. We shall refer to this mechanism as ‘transitional CP violation’. This solution gives a significant hypercharge flux, by the quark or lepton transport[7]. We also showed that a tiny explicit CP violation, which is consistent with the present bound on the neutron EDM, does nonperturbatively resolve the degeneracy between the CP -conjugate pair of the bubbles to leave a sufficient BAU after the EWPT[8].

We examine the possibility of the transitional CP violation at finite temperature in the MSSM by studying the effective potential, when one of the stop mass parameters is small so that the high-temperature expansion is applied only to it.

2 Transitional CP violation

Consider a model with two Higgs doublets whose VEVs are parameterized as

$$\langle \Phi_i \rangle = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \rho_i e^{i\theta_i} \end{pmatrix}, \quad (i = 1, 2) \quad (2.1)$$

and $\theta \equiv \theta_1 - \theta_2$. We assume that the gauge-invariant effective potential near the transition temperature has the form of

$$\begin{aligned} V_{\text{eff}}(\rho_i, \theta_i) &= \frac{1}{2}m_1^2\rho_1^2 + \frac{1}{2}m_2^2\rho_2^2 - m_3^2\rho_1\rho_2 \cos \theta + \frac{\lambda_1}{8}\rho_1^4 + \frac{\lambda_2}{8}\rho_2^4 \\ &+ \frac{\lambda_3 + \lambda_4}{4}\rho_1^2\rho_2^2 + \frac{\lambda_5}{4}\rho_1^2\rho_2^2 \cos 2\theta - \frac{1}{2}(\lambda_6\rho_1^2 + \lambda_7\rho_2^2)\rho_1\rho_2 \cos \theta \\ &- [A\rho_1^3 + \rho_1^2\rho_2(B_0 + B_1 \cos \theta + B_2 \cos 2\theta) \\ &\quad + \rho_1\rho_2^2(C_0 + C_1 \cos \theta + C_2 \cos 2\theta) + D\rho_2^3], \\ &= \left[\frac{\lambda_5}{2}\rho_1^2\rho_2^2 - 2(B_2\rho_1^2\rho_2 + C_2\rho_1\rho_2^2) \right] \\ &\quad \times \left[\cos \theta - \frac{2m_3^2 + \lambda_6\rho_1^2 + \lambda_7\rho_2^2 + 2(B_1\rho_1 + C_1\rho_2)}{2\lambda_5\rho_1\rho_2 - 8(B_2\rho_1 + C_2\rho_2)} \right]^2 \\ &\quad + \theta\text{-independent terms.} \end{aligned} \quad (2.2)$$

The ρ^3 -terms are expected to be induced at finite temperature in a model whose EWPT is of first order. Since we do not consider any explicit CP violation, all the parameters are assumed

to be real. For a given (ρ_1, ρ_2) , the spontaneous CP violation occurs if

$$F(\rho_1, \rho_2) \equiv \frac{\lambda_5}{2} \rho_1^2 \rho_2^2 - 2(B_2 \rho_1^2 \rho_2 + C_2 \rho_1 \rho_2^2) > 0, \quad (2.3)$$

$$-1 < G(\rho_1, \rho_2) \equiv \frac{2m_3^2 + \lambda_6 \rho_1^2 + \lambda_7 \rho_2^2 + 2(B_1 \rho_1 + C_1 \rho_2)}{2\lambda_5 \rho_1 \rho_2 - 8(B_2 \rho_1 + C_2 \rho_2)} < 1. \quad (2.4)$$

In the MSSM at the tree level, $\lambda_5 = \lambda_6 = \lambda_7 = 0$ and $A = B_k = C_k = D = 0$ ($k = 0, 1, 2$), so that no spontaneous CP violation occurs. At zero temperature ($A = B_k = C_k = D = 0$), it is argued that $\lambda_{5,6,7}$ are induced radiatively and (2.3) is satisfied if the contributions from the chargino and neutralino are large enough. For (2.4) to be satisfied, m_3^2 should be as small as $\lambda_6 \rho_1^2 + \lambda_7 \rho_2^2$ so that the pseudoscalar becomes too light[10],

At $T \simeq T_C$, the values of (ρ_1, ρ_2) vary from $(0, 0)$ to $(v_C \sin \beta_C, v_C \cos \beta_C)$ between the symmetric and broken phase regions, where the subscript C denotes the quantities at the transition temperature. Then the effective parameters in (2.2) include the temperature corrections as well. Hence there arises large possibility to satisfy both (2.3) and (2.4) in the intermediate region at the transition temperature, without accompanying too light scalar. If this is the case, a local minimum or a valley of V_{eff} appears for intermediate (ρ_1, ρ_2) with a nontrivial θ . For such a V_{eff} with appropriate effective parameters, the equations of motion for the Higgs fields predict that some class of solutions exist, which have θ of $O(1)$ in the intermediate region even if it vanishes in the broken phase[5].

3 Effective parameters of the MSSM

Since we are concerned with the possibility of the spontaneous CP violation, all the parameters in the lagrangian are assumed to be real. The tree-level Higgs potential of the MSSM is

$$\begin{aligned} V_0 = & m_1^2 \varphi_d^\dagger \varphi_d + m_2^2 \varphi_u^\dagger \varphi_u + (m_3^2 \varphi_u \varphi_d + \text{h.c}) \\ & + \frac{\lambda_1}{2} (\varphi_d^\dagger \varphi_d)^2 + \frac{\lambda_2}{2} (\varphi_u^\dagger \varphi_u)^2 + \lambda_3 (\varphi_u^\dagger \varphi_u) (\varphi_d^\dagger \varphi_d) + \lambda_4 (\varphi_u \varphi_d) (\varphi_u \varphi_d)^* \\ & + \left[\frac{\lambda_5}{2} (\varphi_u \varphi_d)^2 + (\lambda_6 \varphi_d^\dagger \varphi_d + \lambda_7 \varphi_u^\dagger \varphi_u) \varphi_u \varphi_d + \text{h.c} \right], \end{aligned} \quad (3.1)$$

where

$$\begin{aligned} m_1^2 &= \tilde{m}_d^2 + |\mu|^2, & m_2^2 &= \tilde{m}_u^2 + |\mu|^2, & m_3^2 &= \mu B, \\ \lambda_1 &= \lambda_2 = \frac{1}{4}(g_2^2 + g_1^2), & \lambda_3 &= \frac{1}{4}(g_2^2 - g_1^2), & \lambda_4 &= -\frac{1}{2}g_2^2, \end{aligned} \quad (3.2)$$

$$\lambda_5 = \lambda_6 = \lambda_7 = 0, \quad (3.3)$$

Here $g_{2(1)}$ is the $SU(2)(U(1))$ gauge coupling, μ is the coefficient of the Higgs quadratic interaction in the superpotential. The mass squared parameters $\tilde{m}_{u,d}^2$ and μB come from the soft-supersymmetry-breaking terms so that they are arbitrary at this level. m_3^2 could be complex but its phase can be eliminated by the redefinition of the fields when $\lambda_5 = \lambda_6 = \lambda_7 = 0$. We adopt the convention in which this m_3^2 is real and positive.

Let us parameterize the VEVs of the Higgs doublets as

$$\varphi_d = \frac{1}{\sqrt{2}} \begin{pmatrix} \rho_1 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} v_1 \\ 0 \end{pmatrix}, \quad \varphi_u = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \rho_2 e^{i\theta} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 + iv_3 \end{pmatrix}. \quad (3.4)$$

The effective potential at the one-loop level is

$$V_{\text{eff}} = V_0 + V_1(\rho_i, \theta) + \bar{V}_1(\rho_i, \theta; T), \quad (3.5)$$

where $V_1(\rho_i, \theta)$ is the zero-temperature correction given by

$$V_1(\rho_i, \theta) = \sum_j n_j \frac{m_j^4}{64\pi^2} \left[\log \left(\frac{m_j^2}{M_{\text{ren}}^2} \right) - \frac{3}{2} \right], \quad (3.6)$$

and $\bar{V}_1(\rho_i, \theta; T)$ is the finite temperature correction;

$$\bar{V}_1(\rho_i, \theta; T) = \frac{T^4}{2\pi^2} \sum_j n_j \int_0^\infty dx x^2 \log \left[1 - \text{sgn}(n_j) \exp \left(-\sqrt{x^2 + m_j^2/T^2} \right) \right]. \quad (3.7)$$

Here we used the $\overline{\text{DR}}$ -scheme to renormalize V_{eff} with the renormalization scale M_{ren} . n_j counts the degrees of freedom of each species including its statistics, that is, $n_j > 0$ ($n_j < 0$) for bosons (fermions). m_j , which is a function of the Higgs background (ρ_i, θ) , is the mass eigenvalue of each species.

At the one-loop level, $(m_3^2)_{\text{eff}}$ receives corrections only from the Higgs bosons, squarks, sleptons, and charginos and neutralinos. $\lambda_{5,6,7}$, which are zero at the tree level, are generated only through the loops of these particles. Among them, we consider the contributions of charginos($\chi^\pm$), neutralinos(χ^0), stops($\tilde{t}$) and Higgs($\phi^\pm$). The effective parameters are defined as the derivatives of V_{eff} at the origin of the order-parameter space:

$$(m_3^2)_{\text{eff}} = - \left. \frac{\partial^2 V_{\text{eff}}}{\partial v_1 \partial v_2} \right|_0 = m_3^2 + \Delta_\chi m_3^2 + \Delta_{\tilde{t}} m_3^2 + \Delta_{\phi^\pm} m_3^2, \quad (3.8)$$

$$\lambda_5 = \frac{1}{2} \left(\left. \frac{\partial^4 V_{\text{eff}}}{\partial v_1^2 \partial v_2^2} \right|_0 - \left. \frac{\partial^4 V_{\text{eff}}}{\partial v_1^2 \partial v_3^2} \right|_0 \right) = \Delta_\chi \lambda_5 + \Delta_{\tilde{t}} \lambda_5 + \Delta_{\phi^\pm} \lambda_5, \quad (3.9)$$

$$\lambda_6 = -\frac{1}{3} \left. \frac{\partial^4 V_{\text{eff}}}{\partial v_1^3 \partial v_2} \right|_0 = \Delta_\chi \lambda_6 + \Delta_{\tilde{t}} \lambda_6 + \Delta_{\phi^\pm} \lambda_6, \quad (3.10)$$

$$\lambda_7 = -\frac{1}{3} \left. \frac{\partial^4 V_{\text{eff}}}{\partial v_1 \partial v_2^3} \right|_0 = \Delta_\chi \lambda_7 + \Delta_{\tilde{t}} \lambda_7 + \Delta_{\phi^\pm} \lambda_7. \quad (3.11)$$

Here we show only the expressions of the contributions from the charginos, neutralinos and stops, since that from the charged Higgs is much smaller than the others for our choice of the parameters in the next section. As for the derivation, see [11].

When the gaugino mass parameters satisfy $M_2 = M_1$, the contributions from the charginos have the same factors as those from the neutralinos. Then

$$\Delta_\chi m_3^2 = -2g_2^2 \left(1 + \frac{1}{\cos^2 \theta_W} \right) \mu M_2 L(M_2, \mu) + \frac{g_2^2}{\pi^2} \left(1 + \frac{1}{\cos^2 \theta_W} \right) \mu M_2 f_2^{(+)} \left(\frac{M_2}{T}, \frac{\mu}{T} \right),$$

(3.12)

$$\Delta_\chi \lambda_5 = \frac{g_2^4}{8\pi^2} \left(1 + \frac{2}{\cos^4 \theta_W}\right) K\left(\frac{M_2^2}{\mu^2}\right) - \frac{g_2^4}{\pi^2 T^4} \left(1 + \frac{2}{\cos^4 \theta_W}\right) \mu^2 M_2^2 f_4^{(+)}\left(\frac{M_2}{T}, \frac{\mu}{T}\right), \quad (3.13)$$

$$\begin{aligned} \Delta_\chi \lambda_6 &= -\frac{g_2^4}{8\pi^2} \left(1 + \frac{2}{\cos^4 \theta_W}\right) \frac{\mu}{M_2} \left[-H\left(\frac{M_2^2}{\mu^2}\right) + K\left(\frac{M_2^2}{\mu^2}\right) \right] \\ &\quad + \frac{g_2^4}{\pi^2} \left(1 + \frac{2}{\cos^4 \theta_W}\right) \left[\frac{\mu M_2}{T^2} f_3^{(+)}\left(\frac{M_2}{T}, \frac{\mu}{T}\right) + \frac{\mu^3 M_2}{T^4} f_4^{(+)}\left(\frac{M_2}{T}, \frac{\mu}{T}\right) \right] \\ &= \Delta_\chi \lambda_7, \end{aligned} \quad (3.14)$$

where functions $f_n^{(\pm)}(a, b)$, $L(m_1, m_2)$, $K(\alpha)$ and $H(\alpha)$ are defined in [11]. For the condition $F > 0$, it is essential to have a positive λ_5 . The dominant positive contribution comes from (3.13), so that $\mu^2 \simeq M_2^2$ is required since $K(\alpha)$ is maximum at $\alpha = 1$.

The stop contributions are divided into two parts, one of which comes from the light stop, whose supersymmetry-breaking mass parameters are small, while the other originates from the heavier mass eigenstates.

$$\Delta_{\tilde{t}} m_3^2 = N_c y_t^2 \mu A_t L(m_{\tilde{q}}, 0) + \frac{3T^2}{\pi^2} \frac{y_t^2 \mu A_t}{m_{\tilde{q}}^2} \left[I'_B(a_{\tilde{q}}^2) - \frac{\pi^2}{12} \right], \quad (3.15)$$

$$\begin{aligned} \Delta_{\tilde{t}} \lambda_5 &= -\frac{N_c y_t^4}{16\pi^2} \frac{\mu^2 A_t^2}{m_{\tilde{q}}^2 M_{IR}^2} K\left(\frac{m_{\tilde{q}}^2}{M_{IR}^2}\right) \\ &\quad + \frac{N_c y_t^4 \mu^2 A_t^2}{\pi^2 (m_{\tilde{q}}^2)^2} \left[\frac{2T^2}{m_{\tilde{q}}^2} \left(-I'_B(a_{\tilde{q}}^2) + \frac{\pi^2}{12} \right) + I''_B(a_{\tilde{q}}^2) + 2\lambda_- \right], \end{aligned} \quad (3.16)$$

$$\begin{aligned} \Delta_{\tilde{t}} \lambda_6 &= \frac{N_c y_t^2}{16\pi^2} \frac{\mu A_t}{m_{\tilde{q}}^2} \left[\frac{1}{4} \left(\frac{g_1^2}{3} - g_2^2 \right) - \frac{g_1^2 m_{\tilde{q}}^2}{3M_{IR}^2} H\left(\frac{M_{IR}^2}{m_{\tilde{q}}^2}\right) + \frac{y_t^2 \mu^2}{M_{IR}^2} K\left(\frac{m_{\tilde{q}}^2}{M_{IR}^2}\right) \right] \\ &\quad + \frac{N_c y_t^2 \mu A_t}{\pi^2 m_{\tilde{q}}^2} \left\{ \frac{2T^2}{m_{\tilde{q}}^2} \left(\frac{y_t^2 \mu^2}{m_{\tilde{q}}^2} + \left(-\frac{5}{3} g_1^2 + g_2^2 \right) \right) \left[I'_B(a_{\tilde{q}}^2) - \frac{\pi^2}{12} \right] \right. \\ &\quad \left. - \left(\frac{y_t^2 \mu^2}{m_{\tilde{q}}^2} + \frac{3g_2^2 - g_1^2}{12} \right) I''_B(a_{\tilde{q}}^2) + 2 \left(\frac{g_1^2}{3} - \frac{y_t^2 \mu^2}{m_{\tilde{q}}^2} \right) \lambda_- \right\}, \end{aligned} \quad (3.17)$$

$$\begin{aligned} \Delta_{\tilde{t}} \lambda_7 &= \frac{N_c y_t^2}{16\pi^2} \frac{\mu A_t}{m_{\tilde{q}}^2} \left[-\left(y_t^2 + \frac{1}{4} \left(\frac{g_1^2}{3} - g_2^2 \right) \right) \right. \\ &\quad \left. - \left(y_t^2 - \frac{g_1^2}{3} \right) \frac{m_{\tilde{q}}^2}{M_{IR}^2} H\left(\frac{M_{IR}^2}{m_{\tilde{q}}^2}\right) + \frac{y_t^2 A_t^2}{M_{IR}^2} K\left(\frac{m_{\tilde{q}}^2}{M_{IR}^2}\right) \right] \\ &\quad + \frac{N_c y_t^2 \mu A_t}{\pi^2 m_{\tilde{q}}^2} \left\{ \frac{2T^2}{m_{\tilde{q}}^2} \left(\frac{y_t^2 \mu^2}{m_{\tilde{q}}^2} - \left(-\frac{5}{3} g_1^2 + g_2^2 \right) \right) \left[I'_B(a_{\tilde{q}}^2) - \frac{\pi^2}{12} \right] \right. \\ &\quad \left. - \left(y_t^2 + \frac{y_t^2 \mu^2}{m_{\tilde{q}}^2} - \frac{3g_2^2 - g_1^2}{12} \right) I''_B(a_{\tilde{q}}^2) + 2 \left(y_t^2 - \frac{g_1^2}{3} - \frac{y_t^2 \mu^2}{m_{\tilde{q}}^2} \right) \lambda_- \right\}, \end{aligned} \quad (3.18)$$

where $N_c = 3$ and M_{IR} is the infrared cutoff parameter, which will be taken to be the order of the transition temperature. $I_B(a^2)$ is defined in [11].

Table 1: The parameters used in the numerical analysis in the case of $\mu A_t > 0$.

m_3^2	A_t	M_2	μ	$m_{\tilde{g}}$
3300	10	-400	200	400




Figure 1: $(m_3^2)_{\text{eff}}$ and λ_5 as functions of temperature T . The total values are given by the solid curves, the corrections from the stop, chargino-neutralino are depicted by the dashed, dotted and dotted-dashed curves, respectively.

4 Numerical Results

We examine whether the conditions $G(\rho_1, \rho_2) < 1$ and $|G(\rho_1, \rho_2)| < 1$ are satisfied or not by evaluating the effective parameters included in F and G . We take $v_0 = 246\text{GeV}$, $m_t = 177\text{GeV}$, $M_{\text{ren}} = \text{GeV}$ and $\tan\beta_0 = 5$. The mass parameters in the Higgs potential is given by the relations which determine the tree-level minimum of the potential:

$$\begin{aligned} m_1^2 &= m_3^2 \tan\beta_0 - \frac{1}{2}m_Z^2 \cos(2\beta_0), \\ m_2^2 &= m_3^2 \cot\beta_0 + \frac{1}{2}m_Z^2 \cos(2\beta_0). \end{aligned} \quad (4.1)$$

(A) $\mu A_t > 0$

In this case, we take, as the remaining parameters, the values in Table 1. The behaviors of the effective parameters are shown in the Figs. 1 and 2.

The behaviors of the effective parameters are shown in the Figs. 1 and 2. In this case, we have

$$B_2/T = 1.532346 \times 10^{-5}, \quad C_1/T = -4.846485 \times 10^{-3}, \quad (4.2)$$

so that $F(\rho_1, \rho_2) > 0$ is satisfied for $\rho_2 > 5.104$ at $T = 100\text{GeV}$. The region in which $|G(\rho_1, \rho_2)| < 1$ is satisfied are depicted in Figs. 3 and 3.




Figure 2: λ_6 and λ_7 as functions of temperature.




Figure 3: Contour plots of $|G(\rho_1, \rho_2)|$ at $T = 80$ and 90 . $|G(\rho_1, \rho_2)|$ is satisfied in the black region.




Figure 4: The same as Fig. 3 at $T = 100$ and 110 .

Table 2: The parameters used in the numerical analysis in the case of $\mu A_t < 0$.

m_3^2	A_t	M_2	μ	$m_{\tilde{g}}$
2200	10	300	-300	400




Figure 5: $(m_3^2)_{\text{eff}}$ and λ_5 as functions of temperature. The total values are given by the solid curves, the corrections from the stop and chargino-neutralino are depicted by the dashed and dotted curves, respectively.




Figure 6: λ_6 and λ_7 as functions of temperature.




Figure 7: Contour plots of $|G(\rho_1, \rho_2)|$ at $T = 60$ and 70 . $|G(\rho_1, \rho_2)|$ is satisfied in the black region.




Figure 8: The same as Fig. 7 at $T = 80$ and 90 .

(B) $\mu A_t < 0$

The input parameters are given in Table 2. The temperature-dependences of the effective parameters are shown in Figs. 5 and 5. For this parameter set, we have

$$B_2/T = 1.657862 \times 10^{-5}, \quad C_1/T = 3.470457 \times 10^{-3}, \quad (4.3)$$

which implies $F(\rho_1, \rho_2) > 0$ for $\rho_2 > 8.463$ at $T = 100$. The region in which $|G(\rho_1, \rho_2)| < 1$ is satisfied are depicted in Figs. 7 and 8. Note that both λ_6 and λ_7 are negative, while C_1 is negative for the case (A) and positive for (B). This makes the ρ -dependence of $G(\rho_1, \rho_2)$ in the case (B) weaker than in the case (A). Hence The condition $|G(\rho_1, \rho_2)| < 1$ is satisfied in broader region for the case (B).

5 Discussions

The two conditions for the transitional CP violation is satisfied at $T \simeq T_C$, if (i) $|\mu| \simeq |M_{2,1}|$ for $\Delta_\chi \lambda_5 > 0$, (ii) $\mu M_2 < 0$ and m_3^2 to decrease $(m_3^2)_{\text{eff}}$ by the χ -contributions and (iii) m_3^2 is tuned at the tree-level for $(m_3^2)_{\text{eff}}$ to become as small as $\lambda_{6,7}\rho^2$. This mechanism of CP violation

is attractive, since it will be free from the constraints on CP violation at $T = 0$, it can generate sufficient BAU and it is not bothered by the light-scalar problem. Note that for the parameters that admit the transitional CP violation, CP is *not* spontaneously broken at $T = 0$.

It should be noted that the spontaneously- CP -breaking minimum does not have to be the global minimum of V_{eff} . The transitional CP violation could take place if the conditions are satisfied for some fixed (ρ_1, ρ_2) , since such a bubble wall with transitional CP violation would have a lower energy than that without CP violation. For another reason, however, we should understand the global structure of the effective potential, which determines T_C , to know whether the transitional CP violation occurs or not. In this sense, the conditions we examined here should be regarded as the necessary conditions but not the sufficient ones. With the knowledge of the global structure of V_{eff} , one could find the CP -violating profile of the bubble wall so that one could estimate the generated baryon number.

References

- [1] For a review see, A. Cohen, D. Kaplan and A. Nelson, Ann. Rev. Nucl. Part. Sci. **43** (1993) 27.
K. Funakubo, Prog. Theor. Phys. **96** (1996) 475.
- [2] M. Carena, M. Quiros, C.E.M. Wagner, Phys. Lett. **B380** (1996) 81.
D. Delepine, J.-M. Gerard, R. Gonzalez Felipe, J. Weyers, Phys. Lett. **B386** (1996) 183.
J.R. Espinosa, Nucl. Phys. **B475** (1996) 273.
B. de Carlos and J.R. Espinosa, Nucl. Phys. **B503** (1997) 24.
- [3] P. Huet and A. Nelson, Phys. Rev. **D53** (1996) 4578.
M. Aoki, N. Oshimo and A. Sugamoto, Prog. Theor. Phys. **98** (1997) 1179.
M. P. Worah, Phys. Rev. Lett. **79** (1997) 3810.
- [4] A. Nelson, D. Kaplan and A. Cohen, Nucl. Phys. **B373** (1992) 453.
- [5] K. Funakubo, A. Kakuto, S. Otsuki, K. Takenaga and F. Toyoda, Phys. Rev. **D50** (1994) 1105.
- [6] K. Funakubo, A. Kakuto, S. Otsuki, K. Takenaga and F. Toyoda, Prog. Theor. Phys. **94** (1995) 845.
- [7] K. Funakubo, A. Kakuto, S. Otsuki, K. Takenaga and F. Toyoda, Prog. Theor. Phys. **93** (1995) 1067.
- [8] K. Funakubo, A. Kakuto, S. Otsuki and F. Toyoda, Prog. Theor. Phys. **96** (1996) 771.
- [9] K. Funakubo, A. Kakuto, S. Otsuki, F. Toyoda, Prog. Theor. Phys. **98** (1997) 427.
- [10] N. Maekawa, Phys. Lett. **B282** (1992) 387; Nucl. Phys. Suppl. **37A** (1994) 191.
- [11] K. Funakubo, A. Kakuto, S. Otsuki, F. Toyoda, hep-ph/9802276.